arXiv:1410.5867v1 [math.NA] 21 Oct 2014

Solving nonlinear equations by a derivative-free form of the King's family with memory

Somayeh Sharifi[a*] Stefan Siegmund[b†] Mehdi Salimi [c‡]

[a]Young Researchers and Elite Club, Hamedan Branch, Islamic Azad University, Hamedan, Iran
[b,c]Center for Dynamics, Department of Mathematics, Technische Universität Dresden, 01062 Dresden, Germany

Abstract

In this paper, we present an iterative three-point method with memory based on the family of King's methods to solve nonlinear equations. This proposed method has eighth order convergence and costs only four function evaluations per iteration which supports the Kung-Traub conjecture on the optimal order of convergence. An acceleration of the convergence speed is achieved by an appropriate variation of a free parameter in each step. This self accelerator parameter is estimated using Newton's interpolation polynomial of fourth degree. The order of convergence is increased from 8 to 12 without any extra function evaluation. Consequently, this method, possesses a high computational efficiency. Finally, a numerical comparison of the proposed method with related methods shows its effectiveness and performance in high precision computations.

Keywords: Multi-point method, Nonlinear equations, Method with memory, R-order of convergence, Kung-Traub's conjecture.
2010 Mathematics Subject Classification, 65H05.

1 Introduction

Solving nonlinear equations is one of the most important problems that has interesting applications in all fields in science and engineering. We consider iterative methods to find a simple root α of a nonlinear equation $f(x) = 0$, where $f : D \subset \mathbb{R} \to \mathbb{R}$ for an open interval D is a scalar function. Newton-Raphson iteration $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$ is probably the most widely used algorithm for finding roots. It is of second order and requires two evaluations for each iteration step, one evaluation of f and one of f' [12, 18].

*s.sharifi@iauh.ac.ir
†stefan.siegmund@tu-dresden.de
‡Corresponding author: mehdi.salimi@tu-dresden.de

Kung and Traub [8] conjectured that no multi-point method without memory with n evaluations could have a convergence order larger than 2^{n-1}. A multi-point method with convergence order 2^{n-1} is called optimal. The efficiency index, gives a measure of the balance between those quantities, according to the formula $p^{1/n}$, where p is the convergence order of the method and n is the number of function evaluations per iteration.

Some well-known optimal two-point methods have been introduced by Jarratt [6], King [7] and Ostrowski [12]. Some optimal three-point methods have been proposed by Chun and Lee [1], Cordero et al. [2, 3], Lotfi et al. [9], Neta [11], Salimi et al. [15] and Sharma et al. [17]. Zheng et al. [20] have presented an optimal Steffensen-type family and Dzunic et al. [4], Petkovic [13] and Sharma [16] have constructed methods with memory to solve nonlinear equations.

In this paper, we present a modification of the family of King's methods which is derivative-free. The result of this paper is organized as follows: Section 2 and 3 are devoted to the construction and convergence analysis of two-point and three-point optimal derivative-free methods. We introduce the methods with memory and prove their R-order in Section 4. In Section 5, the new methods are compared with a closest competitor in a series of numerical examples. Section 6 contains a short conclusion.

2 Derivative-free two-point method of fourth order

2.1 Description of derivative-free two-point method

We start with King's family of methods which is one of the most important two-point families for solving nonlinear equations [7].

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ x_{n+1} = y_n - \dfrac{f(y_n)}{f'(x_n)} \cdot \dfrac{f(x_n)+\gamma f(y_n)}{f(x_n)+(\gamma-2)f(y_n)}, \quad (n=0,1,\ldots), \quad \gamma \in \mathbb{R}, \end{cases} \tag{2.1}$$

where x_0 is an initial approximation of a simple zero α of f. The main idea is to construct a derivative-free class of two-point methods with optimal order of convergence four. We consider Steffensen-Like's method for the first step and for the second step approximate $f'(x_n)$ by

$$f'(x_n) \approx \frac{f[y_n, w_n]}{G(t_n)},$$

where $w_n = x_n - \beta f(x_n)$, $\beta \neq 0$, $f[y_n, w_n] = \dfrac{f(y_n)-f(w_n)}{y_n - w_n}$, $t_n = \dfrac{f(y_n)}{f(x_n)}$ and G is a real function. Hence, we obtain

$$\begin{cases} y_n = x_n - \dfrac{\beta f(x_n)^2}{f(x_n)-f(w_n)}, \\ x_{n+1} = y_n - \dfrac{f(x_n)+\gamma f(y_n)}{f(x_n)+(\gamma-2)f(y_n)} \cdot \dfrac{f(y_n)}{f[y_n, w_n]} G(t_n). \end{cases} \tag{2.2}$$

2.2 Convergence analysis

We shall state the convergence theorem for the family of methods (2.2).

Theorem 1. *Let $D \subseteq \mathbb{R}$ be an open interval, $f : D \to \mathbb{R}$ four times continuously differentiable and let $\alpha \in D$ be a simple zero of f. If the initial point x_0 is sufficiently close to α, then the method defined by (2.2) converges to α with order four if the weight function $G : \mathbb{R} \to \mathbb{R}$ is continuously differentiable and satisfies the conditions*

$$G(0) = 1, \qquad G'(0) = 2\gamma - 1.$$

Proof. Let $e_n := x_n - \alpha$, $e_{n,w} := w_n - \alpha$, $e_{n,y} := y_n - \alpha$ and $c_n := \dfrac{f^{(n)}(\alpha)}{n!f'(\alpha)}, \quad n = 2, 3, \ldots$.
Using Taylor expansion of f at α and taking into account $f(\alpha) = 0$, we have

$$f(x_n) = f'(\alpha)(e_n + c_2e_n^2 + c_3e_n^3 + c_4e_n^4) + O(e_n^5), \tag{2.3}$$

then

$$e_{n,w} = w_n - \alpha = (1 - \beta f'(\alpha))e_n - \beta f'(\alpha)c_2e_n^2 + O(e_n^3),$$

and

$$f(w_n) = f'(\alpha)(e_{n,w} + c_2e_{n,w}^2 + c_3e_{n,w}^3 + c_4e_{n,w}^4) + O(e_{n,w}^5). \tag{2.4}$$

From (2.3) and (2.4), we have

$$\frac{\beta f(x_n)^2}{f(x_n) - f(w_n)} = e_n - (1 - \beta f'(\alpha))c_2e_n^2 + O(e_n^3), \tag{2.5}$$

by substituting (2.5) in (2.2), we get

$$e_{n,y} = y_n - \alpha = (1 - \beta f'(\alpha))c_2e_n^2 + O(e_n^3).$$

As well as

$$f(y_n) = f'(\alpha)(e_{n,y} + c_2e_{n,y}^2 + c_3e_{n,y}^3 + c_4e_{n,y}^4) + O(e_{n,y}^5). \tag{2.6}$$

From (2.3) and (2.6), we obtain

$$\begin{aligned} t_n = \frac{f(y_n)}{f(x_n)} &= (1 - \beta f'(\alpha))c_2e_n \\ &+ \left(-(3 + \beta f'(\alpha)(-3 + \beta f'(\alpha)))c_2^2 + (-2 + \beta f'(\alpha))(-1 + \beta f'(\alpha))c_3\right)e_n^2 + O(e_n^3), \end{aligned} \tag{2.7}$$

by expanding $G(t_n)$ around 0, we have

$$G(t_n) = G(0) + G'(0)t_n + O(t_n^2), \tag{2.8}$$

moreover, from (2.4) and (2.6), we obtain

$$\begin{aligned} f[y_n, w_n] = \frac{f(y_n) - f(w_n)}{y_n - w_n} &= f'(\alpha) + f'(\alpha)(1 - \beta f'(\alpha))c_2e_n \\ &+ f'(\alpha)((1 - 2\beta f'(\alpha))c_2^2 + (-1 + \beta f'(\alpha))^2c_3)e_n^2 + O(e_n^3), \end{aligned} \tag{2.9}$$

then, by substituting (2.3)-(2.9) in (2.2) we get

$$e_{n+1} = z_n - \alpha = R_2 e_n^2 + R_3 e_n^3 + R_4 e_n^4 + O(e_n^5),$$

with

$$\begin{aligned}
R_2 &= -c_2(1-\beta f'(\alpha))(-1+G(0)),\\
R_3 &= c_2^2(1-\beta f(\alpha))^2(-1+2\gamma-G'(0)),\\
R_4 &= -c_2(1-\beta f'(\alpha))^2(c_2^2(-1-\beta f'(\alpha)-2\gamma^2(1-\beta f'(\alpha)))+c_3).
\end{aligned}$$

Therefore, to provide the fourth order convergence of the two-point method (2.2), it is necessary to choose $R_i = 0$ for $i = 2, 3$, and to achieve this we use the fact that

$$\begin{aligned}
R_2 = 0 \quad &\text{if} \quad G(0) = 1,\\
R_3 = 0 \quad &\text{if} \quad G'(0) = 2\gamma - 1.
\end{aligned}$$

It is clear that $R_4 \neq 0$ in general. Thus, method (2.2) converges to α with order four and the error equation becomes

$$e_{n+1} = -c_2\left(1-\beta f'(\alpha)\right)^2\left(c_2^2(-1-\beta f'(\alpha)-2\gamma^2(1-\beta f'(\alpha)))+c_3\right)e_n^4 + O(e_n^5). \tag{2.10}$$

This finishes the proof of the theorem. □

3 Derivative-free three-point method with order eight

3.1 Description of derivative-free three-point method

Again, we will add one Newton step to the method (2.2)

$$\begin{cases}
y_n = x_n - \dfrac{\beta f(x_n)^2}{f(x_n)-f(w_n)},\\
z_n = y_n - \dfrac{f(x_n)+\gamma f(y_n)}{f(x_n)+(\gamma-2)f(y_n)}\dfrac{f(y_n)}{f[y_n,w_n]}G(t_n),\\
x_{n+1} = z_n - \dfrac{f(z_n)}{f'(z_n)}.
\end{cases} \tag{3.1}$$

As it is seen the function is evaluated for five times, so this method is not optimal and it is also not free of derivatives. To make it an optimal and derivative-free method, we approximate $f'(z_n)$ with Newton's interpolation polynomial of degree three at the point x_n, w_n, y_n and z_n.

$$\begin{aligned}
N_3(t; z_n, y_n, x_n, w_n) := {} & f(z_n) + f[z_n, y_n](t - z_n)\\
& + f[z_n, y_n, x_n](t - z_n)(t - y_n) + f[z_n, y_n, x_n, w_n](t - z_n)(t - y_n)(t - x_n).
\end{aligned}$$

It is clear that

$$N_3(z_n) = f(z_n), \quad \text{and} \quad N_3'(t)\mid_{t=z_n} = f'(z_n).$$

Then

$$N_3'(z_n) = \left[\frac{d}{dt}N_3(t)\right]_{t=z_n} = f[z_n, y_n] + f[z_n, y_n, x_n](z_n - y_n) + f[z_n, y_n, x_n, w_n](z_n - y_n)(z_n - x_n),$$

and hence we get

$$\begin{cases} y_n = x_n - \dfrac{\beta f(x_n)^2}{f(x_n) - f(w_n)}, \\ z_n = y_n - \dfrac{f(x_n) + \gamma f(y_n)}{f(x_n) + (\gamma - 2) f(y_n)} \dfrac{f(y_n)}{f[y_n, w_n]} G(t_n), \\ x_{n+1} = z_n - \dfrac{f(z_n)}{f[z_n, y_n] + f[z_n, y_n, x_n](z_n - y_n) + f[z_n, y_n, x_n, w_n](z_n - y_n)(z_n - x_n)}. \end{cases} \tag{3.2}$$

3.2 Convergence analysis

The following theorem shows that the method (3.2) has convergence order eight.

Theorem 2. *Let $D \subseteq \mathbb{R}$ be an open interval, $f : D \to \mathbb{R}$ eight times continuously differentiable and let $\alpha \in D$ be a simple zero of f. If the initial point x_0 is sufficiently close to α and conditions of Theoerem 1 are established, then the method defined by (3.2) converges to α with order eight.*

Proof. Define $e_n := x_n - \alpha$, $e_{n,w} := w_n - \alpha$, $e_{n,y} := y_n - \alpha$, $e_{n,z} := z_n - \alpha$ and $c_n := \frac{f^{(n)}(\alpha)}{n! f'(\alpha)}$ for $n = 2, 3, \ldots$.

Using Taylor expansion of f at α and taking into account that $f(\alpha) = 0$, we have

$$f(x_n) = f'(\alpha)(e_n + c_2 e_n^2 + c_3 e_n^3 + \ldots + c_8 e_n^8) + O(e_n^9), \tag{3.3}$$

then

$$\begin{aligned} e_{n,w} = w_n - \alpha = (1 - \beta f'(\alpha)) e_n - \beta f'(\alpha) c_2 e_n^2 - \beta f'(\alpha) c_3 e_n^3 - \beta f'(\alpha) c_4 e_n^4 - \beta f'(\alpha) c_5 e_n^5 \\ + O(e_n^6), \end{aligned}$$

and

$$f(w_n) = f'(\alpha)(e_{n,w} + c_2 e_{n,w}^2 + c_3 e_{n,w}^3 + \ldots + c_8 e_{n,w}^8) + O(e_{n,w}^9), \tag{3.4}$$

from (3.3) and (3.4), we have

$$\begin{aligned} e_{n,y} = y_n - \alpha = (1 - \beta f'(\alpha)) c_2 e_n^2 + (-(2 + \beta f'(\alpha)(-2 + \beta f'(\alpha))) c_2^2 + (-2 + \beta f'(\alpha)) \\ (-1 + \beta f'(\alpha)) c_3) e_n^3 + ((4 - \beta f'(\alpha)(5 + \beta f'(\alpha)(-3 + \beta f'(\alpha)))) c_2^3 + (-7 + \beta f'(\alpha) \\ (10 + \beta f'(\alpha)(-7 + 2\beta f'(\alpha)))) c_2 c_3 - (-1 + \beta f'(\alpha))(3 + \beta f'(\alpha)(-3 + \beta f'(\alpha))) c_4) e_n^4 \\ + O(e_n^5), \end{aligned} \tag{3.5}$$

as well as

$$f(y_n) = f'(\alpha)(e_{n,y} + c_2 e_{n,y}^2 + c_3 e_{n,y}^3 + \ldots + c_8 e_{n,y}^8) + O(e_{n,y}^9). \tag{3.6}$$

According to Theorem 1, we get

$$\begin{aligned} e_{n,z} = z_n - \alpha = -c_2 (1 - \beta f'(\alpha))^2 ((-1 - \beta f'(\alpha) + 2\gamma^2 (-1 + \beta f'(\alpha))) c_2^2 + c_3) e_n^4 \\ - (-1 + \beta f'(\alpha))((-4 + \beta f'(\alpha) - (\beta f'(\alpha))^3 - 2\gamma(-1 + \beta f'(\alpha))^3 + 2\gamma^3 \\ (-1 + \beta f'(\alpha))^3 + 2\gamma^2 (-1 + \beta f'(\alpha))(7 + 2\beta f'(\alpha)(-3 + \beta f'(\alpha)))) c_2^4 + (8 - 6\gamma^2 \\ (-2 + \beta f'(\alpha))(-1 + \beta f'(\alpha))^2 + \beta f'(\alpha)(-4 + \beta f'(\alpha)(-5 + 3\beta f'(\alpha)))) c_2 c_3 \\ - (-2 + \beta f'(\alpha))(-1 + \beta f'(\alpha)) c_3^2 - (-2 + \beta f'(\alpha))(-1 + \beta f'(\alpha)) c_2 c_4) e_n^5 + O(e_n^6), \end{aligned}$$

by using the expansion of $f(z_n)$, we have

$$f(z_n) = f^{'}(\alpha)(e_{n,z} + c_2 e_{n,z}^2 + c_3 e_{n,z}^3 + \ldots + c_8 e_{n,z}^8) + O(e_{n,z}^9). \tag{3.7}$$

From (3.6) and (3.7), we get

$$\begin{aligned} f[z_n, y_n] = \frac{f(z_n) - f(y_n)}{z_n - y_n} &= f'(\alpha) + f'(\alpha)c_2^2(1 - \beta f'(\alpha))e_n^2 \\ &+ (-f'(\alpha)(2 + \beta f'(\alpha))c_2^3 + f'(\alpha)(-2 + \beta f'(\alpha))(-1 + \beta f'(\alpha))c_2 c_3)e_n^3 \\ &+ O(e_n^4). \end{aligned} \tag{3.8}$$

And from (3.3) and (3.6), we have

$$\begin{aligned} f[y_n, x_n] = \frac{f(y_n) - f(x_n)}{y_n - x_n} &= f'(\alpha) + c_2 f'(\alpha)e_n + f'(\alpha)(c_2^2(1 - \beta f'(\alpha) + c_3)e_n^2 \\ &+ f'(\alpha)(-(2 + \beta f'(\alpha)(-2 + \beta f'(\alpha)))c_2^3 + (-3 + \beta f'(\alpha))(-1 + \beta f'(\alpha))c_2 c_3 + c_4)e_n^3 \\ &+ O(e_n^4). \end{aligned} \tag{3.9}$$

From (3.3) and (3.4), we obtain

$$\begin{aligned} f[x_n, w_n] = \frac{f(x_n) - f(w_n)}{x_n - w_n} &= f'(\alpha) + c_2 f'(\alpha)(2 - \beta f'(\alpha))e_n + f'(\alpha)(-c_2^2 \beta f'(\alpha) \\ &+ (3 + c_3 \beta f'(\alpha)(-3 + \beta f'(\alpha)))e_n^2 - f'(\alpha)(-2 + \beta f'(\alpha))(-2c_2 c_3 \beta f'(\alpha) \\ &+ (2 + \beta f'(\alpha)(-2 + \beta f'(\alpha)))c_4)e_n^3 + O(e_n^4). \end{aligned} \tag{3.10}$$

From (3.8) and (3.9), we get

$$\begin{aligned} f[z_n, y_n, x_n] = \frac{f[z_n, y_n] - f[y_n, x_n]}{z_n - x_n} &= c_2 f'(\alpha) + c_3 f'(\alpha)e_n + f'(\alpha)(c_2 c_3(1 - \beta f'(\alpha)) + c_4)e_n^2 \\ &+ f'(\alpha)(-c_2^2 c_3(2 + \beta f'(\alpha)(-2 + \beta f'(\alpha))) + c_3^2(-2 + \beta f'(\alpha))(-1 + \beta f'(\alpha)) \\ &+ c_2 c_4(1 - \beta f'(\alpha)) + c_5)e_n^3 + O(e_n^4). \end{aligned} \tag{3.11}$$

And from (3.9) and (3.10), we get

$$\begin{aligned} f[y_n, x_n, w_n] = \frac{f[y_n, x_n] - f[x_n, w_n]}{y_n - w_n} &= c_2 f'(\alpha) + c_3 f'(\alpha)(2 - \beta f'(\alpha))e_n \\ &+ c_2 c_3 f'(\alpha)((1 - 2\beta f'(\alpha)) + (3 + c_4 \beta f'(\alpha)(-3 + \beta f'(\alpha))))e_n^2 \\ &+ O(e_n^3). \end{aligned} \tag{3.12}$$

Therefore, from (3.11) and (3.12), we get

$$\begin{aligned} f[z_n, y_n, x_n, w_n] = \frac{f[z_n, y_n, x_n] - f[y_n, x_n, w_n]}{z_n - w_n} &= c_3 f'(\alpha) + c_4 f'(\alpha)(2 - \beta f'(\alpha))e_n \\ &+ f'(\alpha)(c_2 c_4(1 - 2\beta f(\alpha)) + (3 + c_5 \beta f'(\alpha)(-3 + \beta f'(\alpha)))e_n^2 \\ &+ O(e_n^3). \end{aligned} \tag{3.13}$$

Finally, by substituting (3.3)-(3.13) in (3.2), we obtain

$$\begin{aligned} e_{n+1} = x_{n+1} - \alpha = (1 - \beta f'(\alpha))^4 c_2^2 (c_2^2(-1 - \beta f'(\alpha) - 2\gamma^2(1 - \beta f'(\alpha)) + c_3) \\ (c_2^3(-1 - \beta f'(\alpha) - 2\gamma^2(1 - \beta f'(\alpha))) + c_2 c_3 - c_4)e_n^8 + O(e_n^9). \end{aligned}$$

Which shows that the method (3.2) has optimal convergence order equal to eight. □

4 The development of a new method with memory

In this section, we design a new method with memory by using *self-accelerating parameters* on method (3.2). We observe that the order of convergence of method (3.2) is eight when $\beta \neq 1/f'(\alpha)$. If $\beta = 1/f'(\alpha)$ the convergence order of method (3.2) would be 12. Since the value $f'(\alpha)$ is not available, we use an approximation $\widehat{f}'(\alpha) \approx f'(\alpha)$, instead. The goal is to construct a method with memory that incorporates the calculation of the parameter $\beta = \beta_n$ as the iteration proceeds by the formula $\beta_n = 1/\widehat{f}'(\alpha)$ for $n = 1, 2, 3, \ldots$. It is assumed that an initial estimate β_0 should be chosen before starting the iterative process. In the following, we use the symbols $\to$, O and $\sim$ according to the following convention [18]: If $\lim_{n\to\infty} f(x_n) = C$, we write $f(x_n) \to C$ or $f \to C$, where C is a nonzero constant. If $\frac{f}{g} \to C$, we shall write $f = O(g)$ or $f \sim C(g)$.

We approximate $\widehat{f}'(\alpha)$ by $N_4'(x_n)$, then we have

$$\beta_n = \frac{1}{N_4'(x_n)}, \tag{4.1}$$

where $N_4'(t) := N_4(t; x_n, z_{n-1}, y_{n-1}, w_{n-1}, x_{n-1})$ is Newton's interpolation polynomial of fourth degree, set through five available approximations $(x_n, z_{n-1}, y_{n-1}, w_{n-1}, x_{n-1})$.

$$\begin{aligned} N_4'(x_n) = \left[\frac{d}{dt}N_4(t)\right]_{t=x_n} &= f[x_n, z_{n-1}] + f[x_n, z_{n-1}, y_{n-1}](x_n - z_{n-1}) \\ &+ f[x_n, z_{n-1}, y_{n-1}, w_{n-1}](x_n - z_{n-1})(x_n - y_{n-1}) \\ &+ f[x_n, z_{n-1}, y_{n-1}, w_{n-1}, x_{n-1}](x_n, z_{n-1})(x_n - y_{n-1})(x_n - w_{n-1}). \end{aligned} \tag{4.2}$$

Lemma 3 ([16]). *If $\beta_n = \frac{1}{N_4'(x_n)}$, $n = 1, 2, \ldots$ then the estimate*

$$(1 - \beta_n f'(\alpha)) \sim c_5 e_{n-1} e_{n-1,w} e_{n-1,y} e_{n-1,z}, \tag{4.3}$$

holds.

Theorem 4. *If x_0 is close to a simple zero of the function f, then the convergence R-order of the method (3.2) with memory with the corresponding expression (4.1) of β_n is at least 12.*

Proof. Let (x_n) be a sequence of approximations produced by an iterative method (3.2). If $f(\alpha) = 0$ and this sequence converges to α with the R-order $Q_R((3.2), \alpha) \geq r$, we will write

$$e_{n+1} \sim D_{n,r} e_n^r, \quad e_n = x_n - \alpha, \tag{4.4}$$

where $D_{n,r}$ refers to the asymptotic error of (3.2) when n tend to infinity. In other words, we get

$$e_{n+1} \sim D_{n,r}(D_{n-1,r} e_{n-1}^r)^r = D_{n,r} D_{n-1,r}^r e_{n-1}^{r^2}. \tag{4.5}$$

Assume that the sequences (w_n), (y_n) and (z_n) have the R-order q, p and s, respectively, that is

$$e_{n,w} \sim D_{n,q} e_n^q \sim D_{n,q}(D_{n-1,r} e_{n-1}^r)^q = D_{n,q} D_{n-1,r}^q e_{n-1}^{rq}, \tag{4.6}$$

$$e_{n,y} \sim D_{n,p} e_n^p \sim D_{n,p}(D_{n-1,r} e_{n-1}^r)^p = D_{n,p} D_{n-1,r}^p e_{n-1}^{rp}, \tag{4.7}$$

and

$$e_{n,z} \sim D_{n,s} e_n^s \sim D_{n,s}(D_{n-1,r} e_{n-1}^r)^s = D_{n,s} D_{n-1,r}^s e_{n-1}^{rs}. \tag{4.8}$$

Also, we have

$e_{n,w} \sim (1-\beta_n f'(\alpha))\, e_n,$

$e_{n,y} \sim c_2\,(1-\beta_n f'(\alpha))\, e_n^2,$

$e_{n,z} \sim B\,(1-\beta_n f'(\alpha))^2\, e_n^4,$

where $B = -c_2(c_2^2(-1-\beta_n f'(\alpha)-2\gamma^2(1-\beta_n f'(\alpha)))+c_3),$

$e_{n+1} \sim A(1-\beta_n f'(\alpha))^4 e_n^8,$

where $A = c_2^2(c_2^2(-1-\beta_n f'(\alpha)-2\gamma^2(1-\beta_n f'(\alpha))+c_3)(c_2^3(-1-\beta_n f'(\alpha)-2\gamma^2(1-\beta_n f'(\alpha))+c_2c_3-c_4)$. Therefore by Lemma 3, we obtain

$$\begin{aligned} e_{n,w} &\sim (1-\beta_n f'(\alpha))e_n \sim (c_5 e_{n-1}e_{n-1,w}e_{n-1,y}e_{n-1,z})e_n \\ &\sim c_5 D_{n-1,q}D_{n-1,p}D_{n-1,s}D_{n-1,r}e_{n-1}^{r+p+s+q+1}, \end{aligned} \tag{4.9}$$

$$\begin{aligned} e_{n,y} &\sim c_2(1-\beta_n f'(\alpha))e_n^2 \sim c_2(c_5 e_{n-1}e_{n-1,w}e_{n-1,y}e_{n-1,z})e_n^2 \\ &\sim c_2c_5 D_{n-1,q}D_{n-1,p}D_{n-1,s}D_{n-1,r}^2 e_{n-1}^{2r+s+p+q+1}, \end{aligned} \tag{4.10}$$

$$\begin{aligned} e_{n,z} &\sim B(1-\beta_n f'(\alpha))^2 e_n^4 \sim B(c_5 e_{n-1}e_{n-1,w}e_{n-1,y}e_{n-1,z})^2 e_n^4 \\ &\sim Bc_5^2 D_{n-1,q}D_{n-1,p}^2 D_{n-1,s}^2 D_{n-1,r}^4 e_{n-1}^{4r+2s+2p+2q+2}, \end{aligned} \tag{4.11}$$

$$\begin{aligned} e_{n+1} &\sim A(1-\beta_n f'(\alpha))^4 e_n^8 \sim A(c_5 e_{n-1}e_{n-1,w}e_{n-1,y}e_{n-1,z})^4 e_n^8 \\ &\sim Ac_5^4 D_{n-1,q}D_{n-1,p}^4 D_{n-1,s}^4 D_{n-1,r}^6 e_{n-1}^{8r+4s+4p+4q+4}. \end{aligned} \tag{4.12}$$

By comparing exponents of e_{n-1} appearing in two pairs of relations (4.5), (4.12) and (4.6), (4.9) and (4.7), (4.10) and (4.8), (4.11), we obtain the nonlinear system of four equations and four unknown r, s, p and q.

$$\begin{cases} r^2-8r-4s-4p-4q-4=0, \\ rs-4r-2s-2p-2q-2=0, \\ rp-2r-s-p-q-1=0, \\ rq-r-s-p-q-1=0. \end{cases}$$

A non-trivial solution of the above system is $r=12$, $s=6$, $p=3$, $q=2$.

We have proved that the convergence order of the iterative method (3.2) is at least 12. □

5 Numerical example

In this section, we show the results of some numerical tests to compare the efficiencies of methods, using the programming package Mathematica. To obtain very high accuracy and avoid the loss of significant digits, we employed multi-precision arithmetic with 1200 significant decimal digits in the programming package Mathematica 8.

In what follows, we present some concrete iterative methods from the scheme (3.2).

Method 1. Choose the weight function G as follows:

$$G(t_n) = 1 - t_n, \tag{5.1}$$

where $t_n = \frac{f(y_n)}{f(x_n)}$. The function G in (5.1) satisfies the assumptions of Theorem 2, then we have the following method

$$\begin{cases} y_n = x_n - \dfrac{\beta_n f(x_n)^2}{f(x_n) - f(w_n)}, \quad w_n = x_n - \beta_n f(x_n), \quad \beta_n = \frac{1}{N_4'(x_n)}, \\ z_n = y_n - \dfrac{f(x_n) + \gamma f(y_n)}{f(x_n) + (\gamma - 2) f(y_n)} \cdot \dfrac{f(x_n) - f(y_n)}{f(x_n)} \cdot \dfrac{f(y_n)}{f[y_n, w_n]}, \\ x_{n+1} = z_n - \dfrac{f(z_n)}{f[z_n, y_n] + f[z_n, y_n, x_n](z_n - y_n) + f[z_n, y_n, x_n, w_n](z_n - y_n)(z_n - x_n)}, \end{cases} \tag{5.2}$$

where in general, the divide differential of order n is obtained as:
$f[x_0, x_1, \ldots, x_n] = \dfrac{f[x_1, x_2, \ldots x_n] - f[x_0, x_1, \ldots, x_{n-1}]}{x_n - x_0}$ for $n = 1, 2, \ldots$.

Method 2. Choose the weight function G as follows:

$$G(t_n) = 1 - \frac{t_n}{1 + t_n}, \tag{5.3}$$

where $t_n = \frac{f(y_n)}{f(x_n)}$. The function G in (5.3) satisfies the assumptions of Theorem 2, then we have the following method

$$\begin{cases} y_n = x_n - \dfrac{\beta_n f(x_n)^2}{f(x_n) - f(w_n)}, \quad w_n = x_n - \beta_n f(x_n), \quad \beta_n = \frac{1}{N_4'(x_n)}, \\ z_n = y_n - \dfrac{f(x_n) + \gamma f(y_n)}{f(x_n) + (\gamma - 2) f(y_n)} \cdot \dfrac{f(x_n)}{f(x_n) + f(y_n)} \cdot \dfrac{f(y_n)}{f[y_n, w_n]}, \\ x_{n+1} = z_n - \dfrac{f(z_n)}{f[z_n, y_n] + f[z_n, y_n, x_n](z_n - y_n) + f[z_n, y_n, x_n, w_n](z_n - y_n)(z_n - x_n)}. \end{cases} \tag{5.4}$$

Method 3. Choose the weight function G as follows:

$$G(t_n) = \frac{1 - 2t_n}{1 - t_n}, \tag{5.5}$$

where $t_n = \frac{f(y_n)}{f(x_n)}$. The function G in (5.5) satisfies the assumptions of Theorem 2, then we have the following method

$$\begin{cases} y_n = x_n - \dfrac{\beta_n f(x_n)^2}{f(x_n) - f(w_n)}, \quad w_n = x_n - \beta_n f(x_n), \quad \beta_n = \frac{1}{N_4'(x_n)}, \\ z_n = y_n - \dfrac{f(x_n) + \gamma f(y_n)}{f(x_n) + (\gamma - 2) f(y_n)} \cdot \dfrac{f(x_n) - 2f(y_n)}{f(x_n) - f(y_n)} \cdot \dfrac{f(y_n)}{f[y_n, w_n]}, \\ x_{n+1} = z_n - \dfrac{f(z_n)}{f[z_n, y_n] + f[z_n, y_n, x_n](z_n - y_n) + f[z_n, y_n, x_n, w_n](z_n - y_n)(z_n - x_n)}. \end{cases} \tag{5.6}$$

Method 4. Choose the weight function G as follows:

$$G(t_n) = (1 - t_n)^{\frac{2t_n + 1}{t_n + 1}}, \tag{5.7}$$

where $t_n = \frac{f(y_n)}{f(x_n)}$. The function G in (5.7) satisfies the assumptions of Theorem 2, then we have the following method

$$\begin{cases} y_n = x_n - \dfrac{\beta_n f(x_n)^2}{f(x_n) - f(w_n)}, \quad w_n = x_n - \beta_n f(x_n), \quad \beta_n = \frac{1}{N_4'(x_n)}, \\ z_n = y_n - \dfrac{f(x_n) + \gamma f(y_n)}{f(x_n) + (\gamma - 2) f(y_n)} \cdot \left(\dfrac{f(x_n) - f(y_n)}{f(x_n)} \right)^{\frac{2f(y_n)+f(x_n)}{f(y_n)+f(x_n)}} \cdot \dfrac{f(y_n)}{f[y_n, w_n]}, \\ x_{n+1} = z_n - \dfrac{f(z_n)}{f[z_n, y_n] + f[z_n, y_n, x_n](z_n - y_n) + f[z_n, y_n, x_n, w_n](z_n - y_n)(z_n - x_n)}. \end{cases} \tag{5.8}$$

The new proposed methods with memory were compared with existing three-point methods given in what follows, having the same order convergence and with the same initial data x_0 and β_0.

Method 5. The derivative-free method by Kung and Traub [8] given by

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f[w_n, x_n]}, \quad w_n = x_n + \beta_n f(x_n), \quad \beta_n = \frac{-1}{N'(x_n)}, \\ z_n = y_n - \dfrac{f(y_n) f(w_n)}{(f(w_n) - f(y_n))\, f[x_n, y_n]}, \\ x_{n+1} = z_n - \dfrac{f(y_n) f(w_n) \left(y_n - x_n + \dfrac{f(x_n)}{f[x_n, z_n]} \right)}{(f(y_n) - f(z_n))\, (f(w_n - f(z_n))} + \dfrac{f(y_n)}{f[y_n, z_n]}. \end{cases} \tag{5.9}$$

Method 6. The method by Sharma et al. [16] given by

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{\varphi(x_n)}, \\ z_n = y_n - H(u_n, v_n) \dfrac{f(y_n)}{\varphi(x_n)}, \\ x_{n+1} = z_n - \dfrac{f(z_n)}{f[z_n, y_n] + f[z_n, y_n, x_n](z_n - y_n) + f[z_n, y_n, x_n, w_n](z_n - y_n)(z_n - x_n)}, \end{cases} \tag{5.10}$$

where $\varphi(x_n) = \frac{f(w_n) - f(x_n)}{\beta_n f(x_n)}$, $w_n = x_n + \beta_n f(x_n)$, $\beta_n = \frac{-1}{N'(x_n)}$, $H(u_n, v_n) = \dfrac{1 + u_n}{1 - v_n}$, $u_n = \frac{f(y_n)}{f(x_n)}$ and $v_n = \frac{f(y_n)}{f(w_n)}$.

Method 7. The method by Zheng et al. [20] given by

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f[x_n, w_n]}, \quad w_n = x_n + \beta_n f(x_n), \quad \beta_n = \frac{-1}{N'(x_n)}, \\ z_n = y_n - \dfrac{f(y_n)}{f[y_n, x_n] + f[y_n, x_n, w_n](y_n - x_n)}, \\ x_{n+1} = z_n - \dfrac{f(z_n)}{f[z_n, y_n] + f[z_n, y_n, x_n](z_n - y_n) + f[z_n, y_n, x_n, w_n](z_n - y_n)(z_n - x_n)}, \end{cases} \tag{5.11}$$

In order to test our proposed methods with memory (5.2), (5.4), (5.6) and (5.8) and also compare them with the methods (5.9), (5.10) and (5.11), we choose the initial value x_0 using the `Mathematica` command `FindRoot` [5, pp. 158–160] and compute the error and the computational order of convergence (coc) by the approximate formula [19]

$$\text{coc} \approx \frac{\ln |(x_{n+1} - \alpha)/(x_n - \alpha)|}{\ln |(x_n - \alpha)/(x_{n-1} - \alpha)|}.$$

Test function f_n	Root α
$f_1(x) = \ln(x^2 - 2x + 2) + e^{x^2-5x+4}\sin(x-1)$	1
$f_2(x) = e^{x^2+x\cos(x)-1}\sin(\pi x) + x\ln(x\sin(x)+1)$	0
$f_3(x) = (1-\sin(x^2))\frac{1+x^2}{1+x^3} + x\ln(x^2-\pi+1) - \frac{1+\pi}{1+\sqrt{\pi^3}}$	$\sqrt{\pi}$

Table 1: Test functions f_1, f_2, f_3 and root α.

	Method (5.2)	Method (5.4)	Method (5.6)	Method(5.8)
f_1, $x_0 = 1.35$				
$\lvert x_1 - \alpha\rvert$	$0.314e-6$	$0.134e-5$	$0.543e-6$	$0.703e-6$
$\lvert x_2 - \alpha\rvert$	$0.110e-66$	$0.153e-61$	$0.715e-65$	$0.542e-64$
$\lvert x_3 - \alpha\rvert$	$0.178e-800$	$0.914e-739$	$0.100e-788$	$0.360e-768$
coc	12.1378	12.1056	12.1238	12.1175
f_2, $x_0 = 0.6$				
$\lvert x_1 - \alpha\rvert$	$0.900e-4$	$0.585e-3$	$0.308e-4$	$0.539e-4$
$\lvert x_2 - \alpha\rvert$	$0.111e-44$	$0.713e-38$	$0.650e-49$	$0.335e-48$
$\lvert x_3 - \alpha\rvert$	$0.255e-536$	$0.126e-454$	$0.419e-587$	$0.148e-578$
coc	12.0178	11.9363	12.0465	11.9972
f_3, $x_0 = 1.7$				
$\lvert x_1 - \alpha\rvert$	$0.836e-8$	$0.164e-7$	$0.605e-8$	$0.290e-8$
$\lvert x_2 - \alpha\rvert$	$0.102e-94$	$0.101e-96$	$0.624e-97$	$0.116e-100$
$\lvert x_3 - \alpha\rvert$	$0.134e-1138$	$0.118e-1090$	$0.341e-1165$	$0.612e-1210$
coc	12.0110	12.0173	12.0048	12.0057

Table 2: Errors and coc for methods (5.2), (5.4), (5.6) and (5.8) with $\gamma = 0$.

	Method (5.9)	Method (5.10)	Method (5.11)
f_1, $x_0 = 1.35$			
$\lvert x_1 - \alpha \rvert$	$0.845e-4$	$0.308e-6$	$0.148e-5$
$\lvert x_2 - \alpha \rvert$	$0.393e-45$	$0.179e-67$	$0.157e-61$
$\lvert x_3 - \alpha \rvert$	$0.100e-540$	$0.126e-812$	$0.481e-738$
coc	11.9906	12.1688	12.0973
f_2, $x_0 = 0.6$			
$\lvert x_1 - \alpha \rvert$	$0.798e-3$	$0.891e-4$	$0.214e-4$
$\lvert x_2 - \alpha \rvert$	$0.194e-40$	$0.541e-45$	$0.168e-53$
$\lvert x_3 - \alpha \rvert$	$0.976e-486$	$0.274e-543$	$0.386e-642$
coc	11.8387	12.0827	11.9875
f_3, $x_0 = 1.7$			
$\lvert x_1 - \alpha \rvert$	$0.241e-8$	$0.757e-8$	$0.221e-7$
$\lvert x_2 - \alpha \rvert$	$0.137e-99$	$0.267e-96$	$0.140e-90$
$\lvert x_3 - \alpha \rvert$	$0.283e-1196$	$0.561e-1158$	$0.546e-1089$
coc	12.0190	12.0028	12.0001

Table 3: Errors and coc for methods (5.9), (5.10) and (5.11).

In Table 2, the proposed methods with memory with weight functions (5.1), (5.3), (5.5) and (5.7) and in Tabel 3 the methods (5.9)-(5.11) have been tested on three different nonlinear equations which were introduced in Table 1. It is clear that these methods are in accordance with the developed theory.

6 Conclusion

We have introduced a new method with memory for approximating a simple root of a given nonlinear equation. An increase of the convergence order is attained without any additional function evaluations, which points to a very high computational efficiency of the proposed methods with memory. We have proved that the convergence order of the new method with memory is, at least 12. So its efficiency index is $12^{1/4} = 1.86121$ which is greater than that of the three-point methods of order eight $8^{1/4} = 1.68179$ with the same function evaluations. Numerical examples show that our methods with memory work and can compete with other methods under the same conditions.

References

[1] Chun, C., Lee, M.Y., A new optimal eighth-order family of iterative methods for the solution of nonlinear equations, Appl. Math. Comput., 223, 506-519, (2013).

[2] Cordero, A., Torregrosa, J.R., Vassileva, M.P., Three-step iterative methods with optimal eighth-order convergence, J. Comput. Appl. Math., 235, 3189-3194, (2011).

[3] Cordero, A., Fardi, M., Ghasemi, M., Torregrosa, J.R., Accelerated iterative methods for finding solutions of nonlinear equations and their dynamical behavior, Calcolo, 1-14, (2012).

[4] Dzunic, J., Petkovic, M.S., Petkovic, L.D., Three-point methods with and without memory for solving nonlinear equatins, Appl. Math. Comput., 218, 4917-4927,(2012).

[5] Hazrat, R., Mathematica: A Problem-Centered Approach, Springer-Verlag, 2010.

[6] Jarratt, P., Some fourth order multipoint iterative methods for solving equations, Math. Comput. 20, 434-437, (1966).

[7] King, R.F., Family of four order methods for nonlinear equations SIAM J. Numer. Anal. 10, 876-879, (1973).

[8] Kung, H.T., Traub, J.F., Optimal order of one-point and multipoint iteration, J. Assoc. Comput. Math. 21, 634-651, (1974).

[9] Lotfi, T., Sharifi, S., Salimi, M., Siegmund, S., A new class of three-point methods with optimal convergence order eight and its dynamics, Numer. Algor., DOI 10.1007/s11075-014-9843-y, (2014).

[10] Neta, B., On a family of multipoint methods for nonlinear equations, Intern, J. Computer Math., 9, 353-361, (1981).

[11] Neta, B., On a family of multipoint methods for nonlinear equations, Intern. J. Computer Math., 9, 353-361, (1981).

[12] Ostrowski, A.M., Solution of Equations and Systems of Equations, Academic Press, New York, 1966.

[13] Petkovic, M.S., Dzunic, J., Neta, B., Interpolatory multipoint methods with memory for solving nonlinear equations, Appl. Math. Comput., 218, 2533-2541, (2011).

[14] Petkovic, M.S., Neta, B., Petkovic, L.D., Dzunic, J., Multipoint Methods for Solving Nonlinear Equations, Elsevier, Waltham, MA, 2013.

[15] Salimi, M., Lotfi, T., Sharifi, S., Seigmund, S., Optimal Newton-Secant like methods without memory for solving nonlinear equations with its dynamics, Numer. Algor., Revised, (2014).

[16] Sharma, J.R., Guha, R.K., Gupa, P. Some efficient derivative free methods with memory for solving nonlinear equations, Appl. Math. Comput., 219, 699-707, (2012).

[17] Sharma, J.R., Sharma, R., A new family of modified Ostrowski's methods with accelerated eighth order convergence, Numer. Algor., 54, 445-458, (2010).

[18] Traub, J.F., Iterative Methods for the Solution of Equations, Prentice Hall, New York, 1964.

[19] Weerakoon, S., Fernando, T.G.I., A variant of Newton's method with accelerated third-order convergence, Appl. Math. Lett., 13, 87-93, (2000).

[20] Zheng, Q., Li, J., Huang, F., An optimal Steffensen-type family for solving nonlinear equations, Appl. Math. Comput., 217, 9592-9597, (2011).